PROPOSED REGULATION OF THE

STATE ENVIRONMENTAL COMMISSION

LCB File No. R128-01

November 6, 2001

EXPLANATION - Matter in *italics* is new; matter in brackets formitted material is material to be omitted.

AUTHORITY: §§1-12, NRS 445A.425 and 445A.520.

Section 1. Chapter 445A of NAC is hereby amended by adding thereto a new section to read as follows:

STANDARDS OF WATER QUALITY

East Walker River at Bridge B-1475

Control Point at the East Walker River at Bridge B-1475. The limits of this table apply only from the East Walker River at Bridge B-1475 to the East Walker River at the state line.

PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature Single Value	$\Delta T = 0 \% Ca$	NovApr.: ≤13 °C May-Jun.: ≤17 °C JulOct.: ≤23 °C ΔT ≤2 °Ca	Propagation of aquatic life and recreation involving contact with the water.
pH Single Value		Within range 6.5 - 9.0 SU ∆pH: ±0.5 SU Max.	Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) Annual Average		≤0.10 mg/l	Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and recreation not involving contact with the water.
Nitrogen Species (as N) Annual Average Single Value Single Value	Total Nitrogen ⊴0.9 mg/l ≤1.7mg/l	Nitrate: ≤10 mg/l Nitrite: ≤.06 mg/l	Municipal or domestic supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.

	Ammonia: ≤.02 mg/l (un-ionized)	
	NovMay: ≥6.0 mg/l June-Oct.: ≥5.0 mg/l	Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and recreation not involving contact with the water.
	≤80 mg/l	Propagation of aquatic life.
	b	Propagation of aquatic life and municipal or domestic supply, or both.
	⊴5 PCU	Municipal or domestic supply, or both, and propagation of aquatic life.
≤320 mg/l ≤390 mg/l	≤500 mg/l	Municipal or domestic supply, or both, irrigation and watering of livestock.
≤13 mg/l ≤19 mg/l	∠ 50 mg/l	Municipal or domestic supply, or both, propagation of wildlife, irrigation and watering of livestock.
	≤ 250 mg/l	Municipal or domestic supply, or both.
	⊴8	Irrigation and municipal or domestic supply, or both.
	less than 25% change from natural conditions	Propagation of aquatic life and propagation of wildlife.
	126 MF/100 ml	Recreation involving contact with the water, recreation not involving contact with the water, municipal or domestic supply, or both, irrigation and watering of livestock.
		(un-ionized) NovMay: ≥6.0 mg/l June-Oct.: ≥5.0 mg/l

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. Increase in turbidity must not be more than 10 NTU above natural conditions.
 - **Sec. 2.** NAC 445A.159 is hereby amended to read as follows:
- 445A.159 The standards of water quality for the Walker River from Walker Lake to the state line are prescribed in NAC 445A.160 to 445A.169, inclusive [...], and section 1 of this regulation. The beneficial uses for this area are:
 - 1. Irrigation;
 - 2. Watering of livestock;

- 3. Recreation involving contact with the water;
- 4. Recreation not involving contact with *the* water;
- 5. Industrial supply;
- 6. Municipal or domestic supply, or both;
- 7. Propagation of wildlife; and
- 8. Propagation of aquatic life, and more specifically, the species of major concern are:
- (a) In the West Walker River at the state line, *mountain whitefish*, rainbow trout and brown trout;
- (b) In Topaz Lake, rainbow trout, cutthroat trout, brown trout, kokone salmon and silver salmon;
- (c) In the West Walker River from Wellington to the state line, *mountain whitefish*, rainbow trout and brown trout;
- (d) In the West Walker River from its confluence with the East Walker River to Wellington, brown trout and rainbow trout:
 - (e) In Sweetwater Creek, *mountain whitefish*, brown trout, brook trout and rainbow trout;
- (f) In the East Walker River at the state line, mountain [white fish,] whitefish, rainbow trout and brown trout;
- (g) In the East Walker River from Bridge B-1475 to the state line, mountain whitefish, rainbow trout and brown trout;
- (h) In the East Walker River from its confluence with the West Walker River to [the state line,] Bridge B-1475, brown trout and rainbow trout;
- [(h)] (i) In the Walker River from Weber Reservoir to the confluence of the East Walker River and West Walker River, channel catfish and largemouth bass;

[(i)] (j) In the Walker River from the inlet to Walker Lake to Weber Reservoir, channel catfish, largemouth bass [,] and, from February through June when an adequate flow exists, adult Lahontan cutthroat trout [from April through May,] and adult rainbow [trout from April through June; and

$\frac{(j)}{(j)}$ trout; and

445A.160

- (k) In Desert Creek, brown trout, brook trout and rainbow trout.
- **Sec. 3.** NAC 445A.160 is hereby amended to read as follows:

STANDARDS OF WATER QUALITY

West Walker River

Control Point at the West Walker River at the state line. The limits of this table apply only to the West Walker River at the state line.

		1	T
PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature [°C- Maximum ATa] Single Value	July-Oct.: \leq 22°C $\Delta T = \frac{\{0^{\circ}C\}}{2} \theta Ca$	NovApr.: ≤13°C May-Jun.: ≤17°C JulOct.: ≤23°C ΔT [≤2°C] ≤2 °Ca	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.
pH [Units] Single Value		[S.V.: 7.0 8.3] Within range 6.5 - 9.0 SU ΔpH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering,] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [-mg/l] Annual Average		[A Avg.:] ≤0.1 mg/l	[Aquatic lifeb, water contact recreationb,] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	Total Nitrogen [A Avg.: ≤0.6 S.V.: ≤0.9] 20.6 mg/l 20.9 mg/l	Nitrate : {S.V.:} ≤10 mg/l Nitrite : {S.V.:} ≤.06 mg/l Ammonia : {S.V.:} ≤.02 mg/l (un-ionized)	Municipal or domestic [supplyb, aquatic lifeb water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved		[S.V.: Nov. Apr.: ≥6.0	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering.] Propagation of aquatic

Oxygen [-mg/l] Single Value		May Oct.: ≥5.0] NovMay: ≥6.0 mg/l JunOct: ≥5.0 mg/l	life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [mg/l] Annual Average Single Value	[A Avg.:] ≤60 mg/l	[S.V.:] ≤80 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [-NTU] Single Value		[d] b	[Aquatic Lifeb.] Propagation of aquatic life and municipal or domestic supply [.], or both.
Color [PCU] Single Value	[] ∠ 6 PCU	[e] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [.], or both, and propagation of aquatic life.
Total Dissolved Solids [mg/l] Annual Average Single Value	[A Avg.: ≤165 8.V.: ≤220] ≤165 mg/l ≤20 mg/l	[A Avg.:] ≤500 mg/l	Municipal or domestic [supplyb.] supply, or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average Single Value	[A Avg.: ≤15 S.V.: ≤20] ≤15 mg/l ≤20 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.
Sulfate [-mg/l] Single Value	[] ≤25 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average		[A Avg.:] ≤8	[Irrigationb] Irrigation and municipal or domestic supply [-], or both.
Alkalinity (as CaCO 3) [-mg/l]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
[Fecal Coliform- No./100 ml] Escherichia coli	[A.G.M.: ≤100]	[<u><200/400e</u>]	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or
Annual Geometric Mean Single Value		126 MF/100 ml 235 MF/100 ml	domestic supply, or both, irrigation [, wildlife propagation and stock watering.] and watering of livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. [The most restrictive beneficial use.
- c. Increase in color must not be more than 10 PCU above natural conditions.
- d.] Increase in turbidity must not be more than 10 NTU above natural conditions.
- [e. Based on the minimum of not less than 5 samples taken over a 30 day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30 day period exceed 400 per 100 ml.]
 - **Sec. 4.** NAC 445A.161 is hereby amended to read as follows:

Topaz Lake

Control Point at Topaz Lake. The limits of this table apply at various points in Topaz Lake.

		I	1
PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature [°C- Maximum ATa] Single Value	$\Delta \mathbf{T} = \frac{0^{\circ} \mathbf{C}}{0^{\circ} \mathbf{C}} 0 \mathbf{C} \mathbf{a}$	NovApr.: ≤13°C May-Jun.: ≤17°C JulOct.: ≤23°C ΔT [≤2°C] ≤2 °Ca	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.
pH [Units] Single Value		[S.V.: 7.0 8.3] Within range 6.5 -9.0 SU ΔpH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering.] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [- mg/l] Annual Average Single Value	=	[A Avg.: ≤0.05 — S.V.: ≤0.10] ≤0.05 mg/l ≤0.10 mg/l	[Aquatic lifeb, water contact recreationb.] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	Total Nitrogen [A Avg.: ≤0.6 S.V.: ≤1.0] ≤0.6 mg/l ≤1.0 mg/l	Nitrate: [S.V.:] ≤10 mg/l Nitrite: [S.V.:] ≤.06 mg/l Ammonia: [S.V.:] ≤.02 mg/l (un-ionized)	Municipal or domestic [supplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved Oxygen [-mg/l] Single Value		[S.V.: Nov. Apr.: ≥6.0 May Oct.: ≥5.0] NovMay: ≥6.0 mg/l June-Oct.b: ≥5.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering.] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [-mg/l] Annual Average Single Value	[A Avg.: ≤6.0 S.V.: ≤9.0] \$0.6 mg/l \$9.0 mg/l	[S.V.:] ≤25 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [_NTU] Annual Average Single Value	[A Avg.: ≤3.0 — S.V.: ≤5.0] ≤3.0 NTU ≤5.0 NTU	[d] c	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [-], or both.
Color [-PCU] Single Value	[] ≰I PCU	[c] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [-], or both, and propagation of aquatic life.
Total Dissolved Solids [- mg/l]	[A Avg.: ≤105 — S.V.: ≤120]	[A Avg.:]	Municipal or domestic [supplyb.], or both, irrigation and [stock watering.] watering of livestock.

Annual Average Single Value	≤105 mg/l ≤120 mg/l	≤500 mg/l	
[Chlorides mg/1] Chloride Annual Average Single Value	[A Avg.: ≤7 — S.V.: ≤10] ≤7 mg/l ≤10 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.
Sulfate [- mg/l] Single Value	≤25 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average		[A Avg.:] ≤8	[Irrigationb] Irrigation, and municipal or domestic supply [-], or both.
Alkalinity (as CaCO 3) [-mg/l]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
[Fecal Coliform- No./100 ml] Escherichia coli Annual Geometric	[A.G.M.: ≤25 — S.V.: ≤100]	<u>[≤200/400e]</u>	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or domestic supply, or both, irrigation [, witdlife
Mean Single Value		126 MF/100 ml 235 MF/100 ml	propagation and stock watering.] and watering of livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. The most restrictive beneficial use.
- c. Increase in color must not be more than 10 PCU above natural conditions.
- d. The dissolved oxygen standard from June to October applies only to the epilimnion.
- c. Increase in turbidity must not be more than 10 NTU above natural conditions.
- [e. Based on the minimum of not less than 5 samples taken over a 30-day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30-day period exceed 400 per 100 ml.]
 - **Sec. 5.** NAC 445A.162 is hereby amended to read as follows:

STANDARDS OF WATER QUALITY

West Walker River

Control Point at the West Walker River near Wellington. The limits of this table apply from the West Walker River near Wellington to the West Walker River at the state line.

	1	<u> </u>	T
PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature [°C-Maximum ATa] Single Value	$\Delta \mathbf{T} = \frac{0^{\circ} \mathbf{C}}{0} 0 \mathbf{C} \mathbf{a}$	NovApr.: ≤13°C May-Jun.: ≤17°C JulOct.: ≤23°C ΔT [≤2°C] ≤2 °Ca	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.
pH [Units] Single Value		[S.V.: 7.0 8.3] Within range 6.5 - 9.0 SU ΔpH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering,] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [-mg/l] Annual Average Single Value	[A-Avg.: ≤0.07 S.V.: ≤0.10] ≤0.07 mg/l ≤0.10 mg/l	[A Avg.:] ≤0.1 mg/l	[Aquatic lifeb, water contact recreationb.] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	Total Nitrogen [A Avg.: ≤0.6 S.V.: ≤1.0] ≤0.6 mg/l ≤1.0 mg/l	Nitrate : [S.V.:] ≤10 mg/l Nitrite : [S.V.:] ≤.06 mg/l Ammonia : [S.V.:] ≤.02 mg/l (un-ionized)	Municipal or domestic [supplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved Oxygen [-mg/l] Single Value		[S.V.:] NovMay: ≥6.0 mg/l JunOct.: ≥5.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [-mg/l] Single Value		[S.V.:] ≤80 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [- NTU] Single Value		[d]	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [], or both.
Color [-PCU] Single Value		[e] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [-], or both, and propagation of aquatic life.
Total Dissolved Solids [-mg/l] Annual Average Single Value	[A-Avg.: ≤175 S.V.: ≤260] ≤175 mg/l ≤260 mg/l	[A Avg.:] ≤500 mg/l	Municipal or domestic [supplyb.] supply, or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average Single Value	[A Avg.: ≤16 S.V.: ≤30] ≤16 mg/l ≤30 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.
Sulfate [- mg/l] Single Value		[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average		[A Avg.:] ≤8	[Irrigationb] Irrigation, and municipal or domestic supply [-], or both.
Alkalinity (as CaCO ₃) [- mg/l]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
Fecal Coliform No./100 ml] Escherichia coli Annual Geometric	[A.G.M.: ≤50 — S.V.: ≤150]	[≤200/400e]	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or domestic supply, or both, irrigation [, wildlife]

Mean	126 MF/100 ml	propagation and stock watering.] and watering of
Single Value	 235 MF/100 ml	livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. The most restrictive beneficial use.

- c. Increase in color must not be more than 10 PCU above natural conditions.
- d. Increase in turbidity must not be more than 10 NTU above natural conditions.
- [e. Based on the minimum of not less than 5 samples taken over a 30 day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30 day period exceed 400 per 100 ml.]
 - **Sec. 6.** NAC 445A.163 is hereby amended to read as follows:

STANDARDS OF WATER QUALITY

West Walker River

Control Point at the West Walker River above the confluence with the East Walker River at Nordyke Road. The limits of this table apply to the West Walker River above its confluence with the East Walker River to the control point mentioned in NAC 445A.162 (near Wellington).

PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature [°C Maximum ATa] Single Value	$\Delta T = \begin{bmatrix} 0^{\circ}C \end{bmatrix} 0 Ca$	NovApr.: ≤13°C May-Jun.: ≤17°C JulOct.: ≤23°C ΔT	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.

	1		T
pH [Units] Single Value		[S.V.: 7.0 - 8.3] Within range 6.5 - 9.0 SU ΔpH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering,] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [- mg/l] Annual Average Single Value	[S.V.:] ≤0.15 mg/l	[A Avg::] ≤0.10 mg/l	[Aquatic lifeb, water contact recreationb.] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	Total Nitrogen [A Avg.:≤1.0 S.V.:≤1.2] ≤1.0 mg/l ≤1.2 mg/l	Nitrate : [S.V.:] ≤10 mg/l Nitrite : [S.V.:] ≤.06 mg/l Ammonia : [S.V.:] ≤.02 mg/l (un-ionized)	Municipal or domestic [supplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved Oxygen [- mg/l] Single Value		[S.V.:] NovMay: ≥6.0 mg/l JunOct.: ≥5.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering.] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [mg/l] Single Value		[S.V.:] ≤80 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [- NTU] Single Value		[d] b	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [-], or both.
Color [- PCU] Single Value	[] ≤46 PCU	[c] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [-], or both, and propagation of aquatic life.
Total Dissolved Solids [-mg/l] Annual Average Single Value	[A Avg.: ≤330 S.V.: ≤425] ≤330 mg/l ≤425 mg/l	[A Avg.:] ≤500 mg/l	Municipal or domestic [supplyb.] supply, or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average Single Value	[A Avg:: ≤22 S.V.: ≤28] ≤22 mg/l ≤28 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.
Sulfate [mg/l] Single Value	[] ≤74 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average		[A Avg.:] ≤8	[Irrigationb] Irrigation and municipal or domestic supply [.], or both.
Alkalinity (as CaCO 3) [- mg/1]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
[Fecal Coliform- No./100 ml] Escherichia coli Annual Geometric Mean	[A.G.M.: ≤125 — S.V.: ≤350]	[≤200/400e] 126 MF/100 ml	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or domestic supply, or both, irrigation [, wildlife propagation and stock watering.] and watering of
Single Value		235 MF/100 ml	livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. [The most restrictive beneficial use.

- c. Increase in color must not be more than 10 PCU above natural conditions.
- d.] Increase in turbidity must not be more than 10 NTU above natural conditions.
- [e. Based on the minimum of not less than 5 samples taken over a 30 day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30 day period exceed 400 per 100 ml.]
 - **Sec. 7.** NAC 445A.164 is hereby amended to read as follows:

STANDARDS OF WATER QUALITY

Sweetwater Creek

Control Point at Sweetwater Creek. The limits of this table apply to Sweetwater Creek from its confluence with the East Walker River to the state line.

PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature [°C- Maximum ATa] Single Value	$\Delta \mathbf{T} = \frac{[0^{\circ}\mathbf{C}]}{0} \mathcal{C} \mathbf{a}$	NovApr.: ≤13°C May-Jun.: ≤17°C JulOct.: ≤23°C ΔT [≤2°C] ≤2 °Ca	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.
pH [Units] Single Value		[S.V.: 7.0 8.3] Within range 6.5 - 9.0 SU ΔpH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering,] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [-mg/l] Annual Average		[A Avg.:] ≤0.1 mg/l	[Aquatic lifeb, water contact recreationb,] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	[Total Nitrates A Avg.: ≤0.25 S.V.: ≤0.45] Total Nitrate ≤0.25 mg/l ≤0.45 mg/l	Nitrate : [S.V.:] ≤10 mg/l Nitrite : [S.V.:] ≤.06 mg/l Ammonia : [S.V.:] ≤.02 mg/l (un-ionized)	Municipal or domestic [supplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved		[S.V.:] NovMay: ≥6.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering.] Propagation of aquatic

Oxygen [-mg/l] Single Value		JunOct.: ≥5.0 <i>mg/l</i>	life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [- mg/l] Single Value	[] ≤45 mg/l	[S.V.:] ≤80 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [-NTU] Single Value		[d] b	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [.], or both.
Color [- PCU] Single Value		[e] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic [supplyb.] supply, or both, and propagation of aquatic life.
Total Dissolved Solids [mg/l] Annual Average Single Value	[A-Avg.: ≤220 S.V.: ≤300] ≤220 mg/l ≤300 mg/l	[A Avg.:] ≤500 mg/l	Municipal or domestic [supplyb.] supply, or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average Single Value	[A Avg.: ≤5 S.V.: ≤7] ≤5 mg/l ≤7 mg/l	 [S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.
Sulfate [-mg/l] Single Value		[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average		[A Avg.:] ≤8	[Irrigationb] Irrigation and municipal or domestic supply [], or both.
Alkalinity (as CaCO 3) [- mg/l]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
[Fecal Coliform- No./100 ml] Escherichia coli Annual Geometric Mean Single Value		[≤200/400e] 126 MF/100 ml 235 MF/100 ml	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or domestic supply, or both, irrigation [, wildlife propagation and stock watering.] watering of livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. [The most restrictive beneficial use.
- c. Increase in color must not be more than 10 PCU above natural conditions.
- d. Increase in turbidity must not be more than 10 NTU above natural conditions.
- [e. Based on the minimum of not less than 5 samples taken over a 30-day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30-day period exceed 400 per 100 ml.]
 - **Sec. 8.** NAC 445A.165 is hereby amended to read as follows:

East Walker River

Control Point at the East Walker River at the state line. The limits of this table apply only to the East Walker River at the state line.

		1	1
PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature [°C- Maximum ATa] Single Value	$\Delta \mathbf{T} = \frac{\{0^{\circ} \mathbf{C}\}}{0} \mathbf{C} \mathbf{a}$	NovApr.: ≤13°C May-Jun.: ≤17°C JulOct.: ≤23°C ΔT [≤2°C] ≼2 °Ca	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.
pH [Units] Single Value		[S.V.: 7.0 8.3] Within range 6.5 - 9.0 SU ΔpH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering.] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [-mg/l] Annual Average		[A-Avg.:] ≤0.1 mg/l	[Aquatic lifeb, water contact recreationb.] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	Total Nitrogen [A Avg.: ≤0.8 — S.V.: ≤1.4] ≰0.8 mg/l ≤1.4 mg/l	Nitrate: $[S.V.:] \le 10 \text{ mg/l}$ Nitrite: $[S.V.:] \le .06 \text{ mg/l}$ Ammonia: $[S.V.:] \le .02 \text{ mg/l}$ (un-ionized)	Municipal or domestic [supplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved Oxygen [- mg/l] Single Value		[S.V.:] NovMay: ≥6.0 mg/l JunOct.: ≥5.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering.] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [mg/l] Single Value	[S.V.:] ≤30 mg/l	[S.V.:] ≤80 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [-NTU] Single Value		[d] b	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [-], or both.
Color [- PCU] Single Value		[c] ≰75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [-], or both, and propagation of aquatic life.
Total Dissolved Solids [- mg/l] Annual Average Single Value	[A Avg.: ≤175 — S.V.: ≤210] ≤175 mg/l ≤10 mg/l	[A Avg.:] ≤500 mg/l	Municipal or domestic [supplyb,] supply, or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average	[A Avg.: ≤5 S.V.: ≤7] ≤5 mg/l	 [<u>S.V.:]</u>	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.

Single Value	≤7 mg/l	≤250 <i>mg/l</i>	
Sulfate [mg/l] Single Value	[] ≤26 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [- SAR] Adsorption Ratio Annual Average	[A-Avg.:] ≤2	[A-Avg.:] ≤8	[Irrigationb] Irrigation and municipal or domestic supply [.], or both.
Alkalinity (as CaCO 3) [- mg/l]	-	less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
[Fecal Coliform- No./100 ml] Escherichia coli Annual Geometric	[A.G.M.: ≤20 — S.V.: ≤50]	<u>[≤200/400e]</u>	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, unnicipal or demostic couply or both irrication [, wildlife.
Annual Geometric Mean Single Value		126 MF/100 ml 235 MF/100 ml	domestic supply, or both, irrigation [, wildlife propagation and stock watering.] and watering of livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. [The most restrictive beneficial use.
- c. Increase in color must not be more than 10 PCU above natural conditions.
- d. Increase in turbidity must not be more than 10 NTU above natural conditions.
- [e. Based on the minimum of not less than 5 samples taken over a 30-day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30-day period exceed 400 per 100 ml.]
 - **Sec. 9.** NAC 445A.166 is hereby amended to read as follows:

STANDARDS OF WATER QUALITY

East Walker River

Control Point at the East Walker River south of Yerington above the confluence with the West Walker River (Nordyke Road). The limits of this table apply to the East Walker River south of Yerington above its confluence with the West Walker River to the [state line.] East Walker River at Bridge B-1475.

	REQUIREMENTS TO MAINTAIN EXISTING HIGHER	WATER QUALITY STANDARDS FOR	BENEFICIAL USES As designated in NAC 445A.159
PARAMETER Temperature [°C-	QUALITY	BENEFICIAL USES NovApr.: ≤13°C	(Most stringent use listed first) [Aquatic lifeb and water contact recreation.]
Maximum ΔTa]		May-Jun.: ≤17°C JulOct.: ≤23°C	Propagation of aquatic life and recreation involving contact with the water.
Single Value	$\Delta T = \frac{[0^{\circ}C]}{0} Ca$	ΔT [≤2°C] ≤2 °C a	
pH [Units] Single Value		[S.V.: 7.0 8.3] Within range 6.5 - 9.0 SU ΔpH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering,] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [-mg/l] Annual Average Single Value		[A Avg.: ≤0.16 S.V.: ≤0.39] ≰0.16 mg/l ≤0.39 mg/l	[Aquatic lifeb, water contact recreationb,] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	Total Nitrogen [A Avg.: ≤0.9 S.V.: ≤1.7] \$0.9 mg/l ≤1.7 mg/l	Nitrate : [S.V.:] ≤10 mg/l Nitrite : [S.V.:] ≤.06 mg/l Ammonia : [S.V.:] ≤.02 mg/l	Municipal or domestic [supplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved Oxygen [- mg/l] Single Value		(un-ionized) [S.V.:] NovMay: ≥6.0 mg/l JunOct.: ≥5.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [mg/l] Single Value		[S.V.:] ≤80 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [- NTU] Single Value		[d] <i>b</i>	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [], or both.
Color [PCU] Single Value		[e] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [-], or both, propagation of aquatic life.
Total Dissolved Solids [- mg/l] Annual Average Single Value	[A Avg.: ≤320 S.V.: ≤390] ≤320 mg/l ≤390 mg/l	[A Avg.:] ≤500 mg/l	Municipal or domestic [supplyb.] supply, or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average Single Value	[A Avg.: ≤13 S.V.: ≤19] ≤13 mg/l ≤19 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.
Sulfate [- mg/l] Single Value	[] ≤44 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average		[A Avg.:] ≤8	[Irrigationb] Irrigation and municipal or domestic supply [.], or both.
Alkalinity (as CaCO 3) [mg/l]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
[Fecal Coliform- No./100 ml] Escherichia coli	[A.G.M.: ≤75 — S.V.: ≤350]	<u>[≤200/400e]</u>	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or

Annual	Average		domestic supply, <i>or both</i> , irrigation [, wildlife
Geometr	ric Mean		propagation and stock watering.] and watering of
Single V	'alue	 235 MF/100 ml	livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. [The most restrictive beneficial use.
- c. Increase in color must not be more than 10 PCU above natural conditions.
- d. Increase in turbidity must not be more than 10 NTU above natural conditions.
- [e. Based on the minimum of not less than 5 samples taken over a 30-day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30-day period exceed 400 per 100 ml.]
 - **Sec. 10.** NAC 445A.167 is hereby amended to read as follows: 445A.167

Walker River

Control Point at the Walker River at the inlet to Weber Reservoir. The limits of this table apply to the Walker River from the inlet to Weber Reservoir to the confluence of the West Walker River and the East Walker River.

PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature [°C Maximum		NovMar.: ≤13°C AprJun.: [≤24°C] 23 % JulOct.: ≤28°C	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.
ATa] Single Value	$\Delta T = \frac{[0^{\circ}C]}{0} Ca$	ΔT ≤2°C	

pH [Units] Single Value		[S.V.: 7.0 8.3] Within range 6.5 - 9.0 SU ΔpH: ±0.5 SU Max.	Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering,] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [- mg/l] Annual Average Single Value		[A Avg.: ≤0.26 S.V.: ≤0.40] ≤0.26 mg/l ≤0.40 mg/l	[Aquatic lifeb, water contact recreationb.] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	Total Nitrogen [A-Avg.: ≤1.2 S.V.: ≤1.5] ≤1.2 mg/l ≤1.5 mg/l	Nitrate : [S.V.:] ≤10 mg/l Nitrite : [S.V.:] ≤10 mg/l Ammonia : [S.V.:] ≤.06 mg/l (un-ionized)	Municipal or domestic [supplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved Oxygen [- mg/l] Single Value	 	[S.V.:] NovMay: ≥6.0 mg/l JunOct.: ≥5.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering.] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [[S.V.:] ≤80 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [<u>NTU</u>] Single Value		d	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [-], or both.
Color [- PCU] Single Value		[e] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [-], or both, and propagation of aquatic life.
Total Dissolved Solids [-mg/l] Annual Average Single Value	[A Avg.: ≤400 S.V.: ≤450] ≤400 mg/l ≤450 mg/l	[A Avg.:] ≤500 mg/l	Municipal or domestic [supplyb.], or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average Single Value	[A Avg.: ≤30 S.V.: ≤35] ≤30 mg/l ≤35 mg/l	 [S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife irrigation and [stock watering.] watering of livestock.
Sulfate [-mg/l] Annual Average Single Value	[A Avg.: ≤95 S.V.: ≤110] ≤95 mg/l ≤110 mg/l	[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average	[SAR A Avg.:] ≤3	[A Avg.:] ≤8	[Irrigationb] Irrigation and municipal or domestic supply [.], or both.
Alkalinity (as CaCO ₃) [-mg/l]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
[Fecal Coliform-No./100 mt] Escherichia coli Annual Geometric Mean Single Value	[A.G.M.: ≤100 —S.V.: ≤200]	[≤200/400e] 126 MF/100 ml 235 MF/100 ml	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or domestic supply, or both, irrigation [, wildlife propagation and stock watering.] and watering of livestock.

a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.

- b. [The most restrictive beneficial use.
- c. Increase in color must not be more than 10 PCU above natural conditions.] The temperature beneficial use standard is ≤1 °C from February through June when Lahontan cutthroat are present in the reach from Walker Lake to Weber Reservoir.
- c. The nitrite beneficial use standard is ≤0.06 mg/l from February through June when

 Lahontan cutthroat trout are present in the reach from Walker Lake to the Weber Reservoir.
- d. Increase in turbidity must not be more than 10 NTU above natural conditions.

 [e. Based on the minimum of not less than 5 samples taken over a 30-day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than

10 percent of the total samples taken during any 30-day period exceed 400 per 100 ml.]

Sec. 11. NAC 445A.168 is hereby amended to read as follows: 445A.168

STANDARDS OF WATER QUALITY

Walker River

Control Point at Schurz Bridge. The limits of this table apply from the inlet to Walker Lake to Weber Reservoir.

PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES As designated in NAC 445A.159 (Most stringent use listed first)
Temperature C Maximum	AT - [000] 4 90a	NovMar.: ≤13°C AprJun.: [≤23°C]	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.
ATa] Single Value	$\Delta T = \frac{\text{O}^{\circ} \text{C}}{0} \text{C} a$	ΔT ≤2°C	
pH [Units] Single Value		[S.V.: 7.0 - 8.3] Within range 6.5 - 9.0 SU ApH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering,] watering of livestock, municipal or domestic supply, or both, and industrial

			supply.
Total Phosphates (as P) [- mg/l] Annual Average Single Value		[A Avg: ≤0.17 S.V.: ≤0.23] ≤0.17 mg/ll ≤0.23 mg/	[Aquatic lifeb, water contact recreationb,] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/l] (as N) Annual Average Single Value Single Value Single Value	Total Nitrogen [A Avg.: ≤1.2 S.V.: ≤1.5] ≤1.2 mg/l ≤1.5 mg/l	Nitrate: $\frac{[S.V.:]}{[S.V.:]} \le 10 mg/l$ Nitrite: $\frac{[S.V.:]}{[S.V.:]} \le 10 mg/l$ Ammonia: $\frac{[S.V.:]}{[S.V.:]} \le 000 mg/l$ (un-ionized)	Municipal or domestic Esupplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved Oxygen [- mg/l] Single Value		[S.V.: Nov. Apr.: ≥6.0 May Oct.: ≥5.0] NovMay: ≥6.0 mg/l June-Oct.: ≥5.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [- mg/l] Single Value	[Ann. Avg.:] ≤60 mg/l	[S.V.:] ≤80 mg/l	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [- NTU] Single Value		đ	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [], or both.
Color [- PCU] Single Value		[c] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [-], or both, and propagation of aquatic life.
Total Dissolved Solids [-mg/l] Annual Average Single Value	[A Avg.: ≤390 — S.V.: ≤570] ≤390 mg/l ≤570 mg/l	[A Avg.:] ≤500 mg/l	Municipal or domestic [supplyb.] supply, or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average Single Value	[A Avg.: ≤23 — S.V.: ≤34] ≤23 mg/l ≤34 mg/l	 [S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.
Sulfate [- mg/l] Single Value		[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average	[SAR A Avg.:] ≤3	[A Avg.:] ≤8	[Irrigationb] Irrigation and municipal or domestic supply [-], or both.
Alkalinity (as CaCO ₃) [-mg/l]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
[Fecal Coliform No./100 ml] Escherichia coli Annual Geometric Mean Single Value	[A.G.M.: ≤50 — S.V.: ≤110]	[≤200/400e] 126 MF/100 ml 235 MF/100 ml	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or domestic supply, or both, irrigation [, wildlife propagation and stock watering.] and watering of livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. [The most restrictive beneficial use.

- e. Increase in color must not be more than 10 PCU above natural conditions.] The temperature beneficial use standard is ≤1 °C from February through June when Lahontan cutthroat trout are present.
- c. The nitrite beneficial use standard is ≤0.06 mg/l from February through June when Lahontan cutthroat trout are present.
- d. Increase in turbidity must not be more than 10 NTU above natural conditions.

 [e. Based on the minimum of not less than 5 samples taken over a 30 day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30 day period exceed 400 per 100 ml.]
 - **Sec. 12.** NAC 445A.169 is hereby amended to read as follows: 445A.169

Desert Creek

Control Point at Desert Creek. The limits of this table apply to Desert Creek from its confluence with the West Walker River to the state line.

PARAMETER	REQUIREMENTS TO MAINTAIN EXISTING HIGHER QUALITY	WATER QUALITY STANDARDS FOR BENEFICIAL USES	BENEFICIAL USES Designated in NAC 445A.159 (Most stringent use listed first)
Temperature CC-Maximum ATal Single Value	$\Delta T = \frac{[0 \circ C]}{0} \mathcal{C} a$	NovApr.: ≤13°C May-Jun.: ≤17°C JulOct.: ≤23°C ΔT <mark>{≤2°C} ≤2 °Ca</mark>	[Aquatic lifeb and water contact recreation.] Propagation of aquatic life and recreation involving contact with the water.
pH [Units] Single Value		[S.V.: 7.0 8.3] Within range 6.5 - 9.0 SU ΔpH: ±0.5 SU Max.	[Water contact recreationb, wildlife propagationb,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, irrigation, [stock watering,] watering of livestock, municipal or domestic supply, or both, and industrial supply.
Total Phosphates (as P) [- mg/l] Annual Average		[A Avg.:] ≤0.1 mg/l	[Aquatic lifeb, water contact recreationb.] Propagation of aquatic life, recreation involving contact with the water, municipal or domestic supply,

Single Value	[S.V.:] ≤0.13 mg/l		or both, and [noncontact recreation.] recreation not involving contact with the water.
Nitrogen Species [(N) mg/4] (as N) Annual Average Single Value Single Value Single Value	Total [Nitrates A Avg:: ≤0.20 S.V::≤0.27] Nitrate £0.20 mg/l £0.27 mg/l	Nitrate: [S.V.:] ≤10 mg/l Nitrite: [S.V.:] ≤.06 mg/l Ammonia: [S.V.:] ≤.02 mg/l (un-ionized)	Municipal or domestic [supplyb, aquatic lifeb, water contact recreation, stock watering, wildlife propagation and noncontact recreation.] supply, or both, propagation of aquatic life, recreation involving contact with the water, watering of livestock, propagation of wildlife and recreation not involving contact with the water.
Dissolved Oxygen [- mg/l] Single Value		[S.V.:] NovMay: ≥6.0 mg/l JunOct.: ≥5.0 mg/l	[Aquatic lifeb, water contact recreation, wildlife propagation, stock watering,] Propagation of aquatic life, recreation involving contact with the water, propagation of wildlife, watering of livestock, municipal or domestic supply, or both, and [noncontact recreation.] recreation not involving contact with the water.
Suspended Solids [- mg/l] Single Value		[<u>S.V.:]</u> ≤80 <i>mg/l</i>	[Aquatic lifeb.] Propagation of aquatic life.
Turbidity [- NTU] Single Value		[d] b	[Aquatic lifeb] Propagation of aquatic life and municipal or domestic supply [], or both.
Color [-PCU] Single Value		[c] ≤75 PCU	[Aquatic lifeb and municipal] Municipal or domestic supply [-], or both, and propagation of aquatic life.
Total Dissolved Solids [mg/l] Annual Average Single Value	[A Avg.:≤110 — S.V.:≤130] ≤110 mg/l ≤130 mg/l	[A-Avg.:] ≤500 mg/l	Municipal or domestic [supplyb.], or both, irrigation and [stock watering.] watering of livestock.
[Chlorides mg/l] Chloride Annual Average Single Value	[A Avg.:≤5 — S.V.:≤7] ≤5 mg/l ≤7 mg/l	 [S.V.:] ≤250 mg/l	Municipal or domestic [supplyb, wildlife propagation,] supply, or both, propagation of wildlife, irrigation and [stock watering.] watering of livestock.
Sulfate [-mg/l] Single Value		[S.V.:] ≤250 mg/l	Municipal or domestic [supplyb.] supply, or both.
Sodium [SAR] Adsorption Ratio Annual Average		[A Avg.:] ≤8	[Irrigationb] Irrigation and municipal or domestic supply [-], or both.
Alkalinity (as CaCO 3) [-mg/l]		less than 25% change from natural conditions	[Aquatic lifeb and wildlife propagation.] Propagation of aquatic life and propagation of wildlife.
Fecal Coliform No./100 ml] Escherichia coli Annual Geometric Mean Single Value	[A.G.M.: ≤100 — S.V.: ≤200]	[≤200/400e] 126 MF/100 ml 235 MF/100 ml	[Water contact recreationb, noncontact recreation,] Recreation involving contact with the water, recreation not involving contact with the water, municipal or domestic supply, or both, irrigation [, wildlife propagation and stock watering.] and watering of livestock.

- a. Maximum allowable increase in temperature above water temperature at the boundary of an approved mixing zone, but the increase must not cause a violation of the single value standard.
- b. The most restrictive beneficial use.
- c. Increase in color must not be more than 10 PCU above natural conditions.
- d. Increase in turbidity must not be more than 10 NTU above natural conditions.

[e. Based on the minimum of not less than 5 samples taken over a 30-day period, the fecal coliform bacterial level may not exceed a geometric mean of 200 per 100 ml nor may more than 10 percent of the total samples taken during any 30-day period exceed 400 per 100 ml.]